

BFV QUANTIZATION OF RELATIVISTIC SPINNING PARTICLES WITH A SINGLE BOSONIC CONSTRAINT

Silvio J. Rabello* and Arvind N. Vaidya

Instituto de Física
Universidade Federal do Rio de Janeiro
Rio de Janeiro RJ
Caixa Postal 68.528-CEP 21945-970
Brasil

February 1, 2008

Abstract

Using the BFV approach we quantize a pseudoclassical model of the spin one half relativistic particle that contains a single bosonic constraint, contrary to the usual locally supersymmetric models that display first and second class constraints.

ℒ_TEX

*e-mail: goldegol@vms1.nce.ufrj.br

The dynamics of the relativistic spinless particle, of mass m and 4-momentum P^μ is generated by the vanishing canonical Hamiltonian $H_0 = P^2 - m^2 \approx 0$ [1], reflecting the invariance under reparametrizations of the world line $X^\mu(s)$ ($s \in [0, 1]$). To describe spinning relativistic particles one often promotes the reparametrization invariance of the spinless action to a local supersymmetry on s [2] (one dimensional supergravity). The resulting pseudoclassical system has first and second class constraints that after an appropriate treatment (e.g. the Dirac algorithm) reduce to two first class constraints, one bosonic the other fermionic, the mass-shell condition and its “square root” respectively. In this paper we study another description of a spinning particle, where only one bosonic constraint is needed. The inspiration for this model comes from the original Dirac Hamiltonian [3] ¹:

$$H_D = \gamma_0 \boldsymbol{\gamma} \cdot \mathbf{P} + \gamma_0 m \quad (1)$$

where $\mathbf{P}$ is the 3-momentum and the γ 's are the Dirac matrices. The above H_D generates translations in the physical time X^0 and it is not a Lorentz scalar. In order to have a model for the relativistic spinning particle in the same lines of the spinless case we study an analog of (1) that is Lorentz covariant, generate translations in the parameter time s and is weakly zero so that the resulting theory is invariant under reparametrizations of $X^\mu(s)$. Introducing the commuting Lagrange multiplier λ and anticommuting coordinates ζ_5 and ζ_μ ($\mu = 0, 1, 2, 3$), the pseudoclassical Lagrangian reads:

$$\mathcal{L} = P^\mu \dot{X}_\mu + \frac{i}{2}(\zeta_5 \dot{\zeta}_5 - \zeta^\mu \dot{\zeta}_\mu) - \lambda H_0, \quad (2)$$

with the canonical Hamiltonian given by

$$H_0 = \zeta_5 \zeta_\mu P^\mu - m \approx 0, \quad (3)$$

From this Lagrangian we can read the symplectic structure that on quantization gives the (anti)commutation relations (the non zero part of it) :

$$[X^\mu, P^\nu]_- = i\eta^{\mu\nu}, \quad [\zeta_\mu, \zeta_\nu]_+ = -2\eta_{\mu\nu}, \quad [\zeta_5, \zeta_5]_+ = 2. \quad (4)$$

¹ $\hbar = c = 1$ and $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$

In the quantum mechanics of this model the constraint (3) becomes a gauge generator, that in the Dirac theory of constrained systems [4] annihilates the physical states, i.e. $H_0|phys\rangle = 0$. For consistency we have that its square,

$$H_0^2 = P^2 + m^2 - 2m\zeta_5\zeta_\mu P^\mu, \quad (5)$$

must also annihilate $|phys\rangle$. It is easy to see that on physical states we have :

$$H_0^2|phys\rangle = (P^2 - m^2)|phys\rangle = 0. \quad (6)$$

Thus our single bosonic constraint (3) implies the mass-shell condition of a relativistic particle of mass m . We now proceed to the BFM quantization of the model.

In the BFM formulation [5, 4] the hermitian nilpotent BRST charge operator for this model is given by

$$Q = \eta H_0 + \bar{\mathcal{P}}\pi, \quad (7)$$

where we extended our phase space by introducing the conjugate momentum π for the Lagrange multiplier λ , a fermionic pair of conjugate ghost operators $(\eta, \mathcal{P})$ for the constraint H_0 , and a pair of fermionic antighost conjugate operators $(\bar{\eta}, \bar{\mathcal{P}})$ to take care of the constraint π that appear due to the “einbein” character of λ that we assume from start. These hermitian operators generate the algebra (the non zero part of it)

$$[\lambda, \pi]_- = i, \quad [\eta, \mathcal{P}]_+ = [\bar{\eta}, \bar{\mathcal{P}}]_+ = 1. \quad (8)$$

Due to the nilpotency of Q the following extension of the original BRST invariant Hamiltonian H_0 is also invariant :

$$H_0 \rightarrow H_0 + [\Psi, Q]_+. \quad (9)$$

Where Ψ is an arbitrary “gauge fermion”, that we here choose to be $\Psi = \mathcal{P}\lambda$. And also, since $Q|phys\rangle = 0$, the above extension of H_0 can be used in the evolution operator without changing the transition amplitudes. We now consider the transition amplitude

$$K(z'', z'; s) \equiv \langle z'' | e^{-iHs} | z' \rangle = \langle z'', s | z', 0 \rangle, \quad (10)$$

in the basis $|z'; s\rangle \equiv |P', \lambda', \eta', \bar{\eta}'; s\rangle$ with $\langle z'', 0|z', 0\rangle = \delta(z'' - z')$ (hereafter all operator eigenvalues will be primed). The above $K(z'', z'; s)$ obeys the Schrödinger equation in the parameter time s :

$$i \frac{\partial K(z'', z'; s)}{\partial s} = \langle z'', s|H|z', 0\rangle, \quad (11)$$

with H being

$$H = [\Psi, Q]_+ = \lambda(\zeta_5 \zeta_\mu P^\mu - m) + i\mathcal{P}\bar{\mathcal{P}}. \quad (12)$$

The expectation value $\langle z'', s|H|z', 0\rangle$ is easily evaluated if we apply a method developed long ago by Schwinger [6] that consists in solving the Heisenberg equations for the canonical variables, writing the $\mathcal{P}$ and $\bar{\mathcal{P}}$ operators in terms of the η and $\bar{\eta}$ operators and inserting these in a time ordered fashion in H . After that we can easily integrate (11). For this model we have the Heisenberg equations (the dot means derivative with respect to s) :

$$\dot{X}^\mu = \lambda \zeta_5 \zeta^\mu, \quad \dot{P}^\mu = 0, \quad (13)$$

$$\dot{\zeta}^\mu = -2i\lambda \zeta_5 P^\mu, \quad \dot{\zeta}_5 = -2i\lambda \zeta_\mu P^\mu, \quad (14)$$

$$\dot{\lambda} = 0, \quad \dot{\pi} = -(\zeta_5 \zeta_\mu P^\mu - m), \quad (15)$$

$$\dot{\eta} = \bar{\mathcal{P}}, \quad \dot{\bar{\mathcal{P}}} = 0, \quad (16)$$

$$\dot{\bar{\eta}} = -\mathcal{P}, \quad \dot{\mathcal{P}} = 0. \quad (17)$$

As in the case of the spinless particle [1] the above equations carry an ambiguity, take the expectation value of $\dot{\zeta}_\mu$, if we change $\lambda \rightarrow -\lambda$ and $P_\mu \rightarrow -P_\mu$ we return to the same equation. A remedy to solve this ambiguity in expectation values, is to restrain the eigenvalues of λ to be either positive or negative [1].

The solutions for the ghosts and antighosts are:

$$\eta(s) = \eta(0) + \bar{\mathcal{P}}s, \quad \bar{\eta}(s) = \bar{\eta}(0) - \mathcal{P}s. \quad (18)$$

We now write $\mathcal{P}$ and $\bar{\mathcal{P}}$ in terms of $\eta(s)$, $\eta(0)$, $\bar{\eta}(s)$ and $\bar{\eta}(0)$. For the ζ 's we take as initial values $\zeta_5(0) = \gamma_5$ and $\zeta_5(0)\zeta_\mu(0) = \gamma_\mu$. Since H is s -independent we from now

on substitute these values in it, so that we get the time ordered Hamiltonian :

$$H_{ord} = \lambda(\gamma_\mu P^\mu - m) - \frac{i}{s^2} \left(\bar{\eta}(s)\eta(s) - \bar{\eta}(s)\eta(0) \right. \\ \left. + \bar{\eta}(0)\eta(0) + \eta(s)\bar{\eta}(0) - [\bar{\eta}(0), \eta(s)]_+ \right), \quad (19)$$

with $[\bar{\eta}(0), \eta(s)]_+ = s$. We are now in position to integrate (11):

$$\ln K = -is\lambda'(\gamma^\mu P'_\mu - m) - \frac{1}{s}(\bar{\eta}'' - \bar{\eta}')(\eta'' - \eta') - \ln s + \Phi, \quad (20)$$

where Φ is a s independent function of the dynamical variables. With aid of the boundary condition $\langle z'', 0 | z', 0 \rangle = \delta(z'' - z')$:

$$\Phi = \ln i \delta^4(P'' - P') \delta(\lambda'' - \lambda'). \quad (21)$$

Now that we have found K we must impose the invariance under Q , i.e. Q acting each argument of $K(z'', z'; s)$ must produce zero as a result. Among the several ways for this to be true [4], we choose the boundary conditions :

$$\pi'' = \pi' = \eta'' = \eta' = \bar{\eta}'' = \bar{\eta}' = 0. \quad (22)$$

The condition on $\bar{\eta}$ is a consistency one for $\pi = [Q, \bar{\eta}]_+$ and Q to annihilate $|phys\rangle$. We now Fourier transform K on λ'' and λ' choosing $\lambda \geq 0$,

$$K = i \int_0^\infty d\lambda' s e^{i\lambda'(\pi'' - \pi')} e^{-i[s\lambda'(\gamma^\mu P'_\mu - m) + \frac{i}{s}(\bar{\eta}'' - \bar{\eta}')(\eta'' - \eta')]} \delta^4(P'' - P'). \quad (23)$$

With the BRST invariant boundary conditions and defining $s\lambda' \equiv T$:

$$K = i \int_0^\infty dT e^{-iT(\gamma^\mu P'_\mu - m)} \delta^4(P'' - P') = \frac{\delta^4(P'' - P')}{\gamma^\mu P'_\mu - m - i0_+}. \quad (24)$$

The choice of positive λ led to the momentum space Feynman propagator for Dirac fermions, had we chosen negative values for λ we would end up with the complex conjugate of it.

Discussion

It would be interesting to introduce interactions with background fields as well as to generalize our Lagrangian to describe extended objects like spinning strings and membranes. This work was partially supported by the CNPq.

References

- [1] M. Henneaux and C. Teitelboim *Ann. Phys.* **143**, 127 (1982).
- [2] L. Brink , P. Di Vecchia and P. Howe *Nucl. Phys. B* **118**, 76 (1977).
- [3] P.A.M. Dirac *The Principles of Quantum Mechanics* (3rd edn) (Oxford University Press, 1947).
- [4] M. Henneaux and C. Teitelboim *Quantization of Gauge Systems* (Princeton Univ. Press, 1992); M. Henneaux *Phys. Rep.* **126**, 1 (1985).
- [5] E.S. Fradkin and G.A. Vilkovisky *Phys. Lett.* **55B**, 224 (1975); I.A. Batalin and G.A. Vilkovisky *Phys. Lett.* **69B**, 309 (1977).
- [6] J. Schwinger *Phys. Rev.* **82**, 664 (1951).